

CLOSED INCOMPRESSIBLE SURFACES IN THE COMPLEMENTS OF POSITIVE KNOTS

MAKOTO OZAWA

ABSTRACT. We show that any closed incompressible surface in the complement of a positive knot is algebraically non-split from the knot, positive knots cannot bound non-free incompressible Seifert surfaces and that the splitability and the primeness of positive knots and links can be seen from their positive diagrams.

1. INTRODUCTION

A knot K in the 3-sphere S^3 is called *positive* if it has an oriented diagram all crossings of which are positive crossings. For a closed surface F in $S^3 - K$, we define the *order* $o(F; K)$ of F for K as follows ([2]). Let $i : F \rightarrow S^3 - K$ be the inclusion map and let $i_* : H_1(F) \rightarrow H_1(S^3 - K)$ be the induced homomorphism. Since $\text{Im}(i_*)$ is a subgroup of $H_1(S^3 - K) = \mathbb{Z}\langle \text{meridian} \rangle$, there is an integer m such that $\text{Im}(i_*) = m\mathbb{Z}$. Then we define $o(S; K) = m$.

The positive knot complements have the following special properties.

Theorem 1.1. *Any closed incompressible surface in a positive knot complement has non-zero order.*

A Seifert surface F for a knot is said to be *free* if $\pi_1(S^3 - F)$ is a free group. In [2, Theorem 1.1], it is shown that a knot bounds a non-free incompressible Seifert surface if and only if there exists a closed incompressible surface in the knot complement whose order is equal to zero. Therefore, Theorem 1.1 gives us the next corollary.

Corollary 1.2. *Positive knots cannot bound non-free incompressible Seifert surfaces.*

Although positive links which have connected positive diagrams are non-split because they have positive linking numbers, we can give another geometrical proof of this fact.

Theorem 1.3. *Positive links are non-split if their positive diagrams are connected.*

Positive diagrams of positive knots or links also tell us their primeness. We say that a knot or link diagram \tilde{K} on the 2-sphere S is *prime* if for any loop l in S intersecting \tilde{K} in 2 points, l bounds a disk intersecting \tilde{K} in an arc.

Theorem 1.4. *Non-trivial positive knots or links are prime if their positive diagrams are connected and prime.*

1991 *Mathematics Subject Classification.* 57M25.

Key words and phrases. positive knot, closed incompressible surface, order, free Seifert surface, splitability, primeness.

The author was supported in part by Fellowship of the Japan Society for the Promotion of Science for Japanese Junior Scientists.

Theorem 1.4 widens the result of Cromwell ([1, 1.2 Theorem]).

2. PROOF OF THEOREM 1.1 AND 1.3

Theorem 1.1 and 1.3 follow the next Theorem.

Theorem 2.1. *Let K be a positive knot or link in the 3-sphere S^3 and F a closed incompressible surface in the complement of K . Then one of the following conclusions (1) and (2) holds.*

- (1) *There exists a loop l in F such that $lk(l, K) \neq 0$.*
- (2) *F is a splitting sphere for K , and any positive diagram of K is disconnected.*

Henceforth, we shall prove Theorem 2.1.

Let S be a 2-sphere in S^3 and $p : S^3 - \{2 \text{ points}\} \cong S \times R \rightarrow S$ a projection. Put K so that $p(K)$ is a positive diagram. As usual way, we express K in a bridge presentation. Thus we have the following data (see Figure 1).

- $S^3 = B^+ \cup_S B^-$ (S decomposes S^3 into two 3-balls)
- $K = K^+ \cup_S K^-$, where $K^\pm \subset B^\pm$ (S cuts K into over bridges and under bridges)
- $K^\pm = K_1^\pm \cup K_2^\pm \cup \dots K_n^\pm$ (K is presented as n over bridges and n under bridges)
- $D^\pm = D_1^\pm \cup D_2^\pm \cup \dots D_n^\pm$ (each $K_i^\pm \cup p(K_i^\pm)$ bounds a disk D_i^\pm such that $p(D_i^\pm) = p(K_i^\pm)$)




FIGURE 1. View from level surface

We take n minimal over all bridge presentations of $p(K)$.

Lemma 2.2. *We may assume that $F \cap D^- = \emptyset$, $F \cap B^-$ consists of disks, $F \cap D^+$ consists of arcs, and any component of $F \cap B^+ - D^+$ is a disk.*

Proof. This can be done by cutting and pasting F along some disks. Note that such operations do not have any effect on the conditions (1) and (2) if we take a suitable choice of F . \square

We take $|F \cap B^-|$ and $|F \cap D^+|$ minimal. Note that $|F \cap B^-| \neq 0$ because F is incompressible in $S^3 - K$. If $|F \cap B^-| = 1$ and $|F \cap D^+| = 0$, then we have the conclusion (2).

Hereafter, we suppose that $|F \cap B^-| \geq 1$ and $|F \cap D^+| \geq 1$.

Then we obtain a connected graph G in F by regarding $F \cap B^-$ and $F \cap D^+$ as vertices and edges respectively. Note that every vertex has a positive even valency by the construction.

An arc α_j of $F \cap D_i^+$ divides D_i^+ into two disks δ_j and δ'_j , where δ'_j contains K_i^+ . Put $\beta_j = \delta_j \cap S$. We may assume that $p(\alpha_j) = p(\delta_j) = \beta_j$ for all α_j . We assign an orientation endowed from K_i to α_j and β_j naturally (see Figure 2).




FIGURE 2. α_j and β_j have the orientations

Lemma 2.3. *For any arc α_j of $F \cap D_i^+$, $\beta_j \cap p(K^-) \neq \emptyset$.*

Proof. Suppose that there exists an arc α_j of $F \cap D_i^+$ such that $\beta_j \cap p(K^-) = \emptyset$. By exchanging α_j if necessary, we may assume that α_j is outermost in D_i^+ , that is, $\text{int}\delta_j \cap F = \emptyset$. If α_j connects different vertices, then a ∂ -compression of F along δ_j reduces $|F \cap B^-|$. Otherwise, α_j incident to a single vertex, say D_k^- . We perform a ∂ -compression of F along δ_j , and obtain an annulus A consisting of the disk D_k^- and the resultant band b . Since we chose an outermost arc α_j and $\beta_j \cap p(K^-) = \emptyset$, there exists a compressing disk for A in $B^- - K^-$. By retaking F along the compressing disk, we can reduce $|F \cap D^+|$. \square

Now we pay attention to a face f of G in F . The ‘cycle’ ∂f consists of edges and ‘corners’ as subarcs in $\partial(F \cap B^-)$. The edges have orientations as previously mentioned.

Lemma 2.4. *For any face f , the cycle ∂f can not be oriented.*

Proof. Suppose that there is a face f such that ∂f can be oriented. Then, since no corner of ∂f intersects $p(K)$, and by Lemma 2.3, $p(\partial f)$ has non-zero intersection number with $p(K^-)$ on S as illustrated in Figure 3. This is a contradiction. \square




FIGURE 3. $p(\partial f)$ has non-zero intersection number

For each face f of G and any point in the interior of any edge of ∂f , we can find an arc γ on f satisfying the following property.

(*) γ connects two edges of ∂f whose orientations are different in ∂f .




FIGURE 4. γ with the property (*)

Lemma 2.4 assures the existence of such an arc γ .

To find a loop l on F with $lk(l, K) \neq 0$, we depart a point in the interior of any edge of G , trace arcs with the property (*), and will arrive at the face on which we have walked. Connecting these arcs, we will obtain an oriented loop l in $F \cap B^+$ with a suitable orientation such that l has a positive intersection number with edges of G on F . Thus we got an oriented loop l in F which has non-zero linking number with K . Since any loop in a splitting sphere is contractible in $S^3 - K$, we have the conclusion (1).

This completes the proof of Theorem 2.1.

3. PROOF OF THEOREM 1.4

Let K be a positive knot or link in S^3 and F be a decomposing sphere for K . We put K and F as the proof of Theorem 1.1 except that two points p_1 and p_2 of $F \cap K$ are in $intB^+$ or $intB^-$. Note that p_1 and p_2 can not be the ends of a single arc of $F \cap D^\pm$ because the tangle (B^\pm, K^\pm) is trivial and F is a decomposing sphere. Hence, there are two arcs e_1 and e_2 of $F \cap D^\pm$ whose ends contain p_1 and p_2 respectively. We deform F by an isotopy relative to K so that $p(e_i) = p(p_i)$ ($i = 1, 2$). We take the number of bridges n minimal.

Lemma 3.1. *We may assume that $F \cap D^- \subset e_1 \cup e_2$, $F \cap B^-$ consists of disks, $F \cap D^+$ consists of arcs, and any component of $F \cap B^+ - D^+$ is a disk.*

Proof. This can be done by an isotopy of F since Theorem 1.3 assures us that $S^3 - K$ is irreducible. \square

We take $|F \cap B^-|$ and $|(F \cap D^+) - (e_1 \cup e_2)|$ minimal. Then we obtain a connected graph G in F by regarding $F \cap B^-$ and $(F \cap D^+) - (e_1 \cup e_2)$ as vertices and edges respectively. Corners of each face of G may contain two points $\partial e_1 - p_1$ and $\partial e_2 - p_2$. Note that $|F \cap B^-| \neq 0$, otherwise F is not a decomposing sphere since (B^\pm, K^\pm)

is a trivial tangle. If $|F \cap B^-| = 1$ and $F \cap D^+ \subset e_1 \cup e_2$, then $F \cap S$ gives a desired loop since $p(e_i) = p(p_i)$ ($i = 1, 2$).

Lemma 3.2. *For any arc α_j of $(F \cap D^+) - (e_1 \cup e_2)$, $\beta_j \cap p(K^-) \neq \emptyset$.*

Proof. This can be done by the same argument to Lemma 2.3. \square

Hereafter, we assume that \tilde{K} is prime.

Lemma 3.3. *There is no vertex of G with valency 1.*

Proof. Suppose that there is a vertex V with valency 1. Then only one edge α incident to V , and hence exactly one of e_1 and e_2 is attached to V or contained in V . Thus ∂V intersects \tilde{K} in two points. Since \tilde{K} is prime, ∂V bounds a disk E in S which intersects $p(K)$ in an unknotted arc. In the former case, $p(K) \cap E$ lies under a subarc of K^+ by the minimality of the number of bridges n . Then by an isotopy of F along the 3-ball which is bounded by $V \cup E$, we can reduce $|F \cap B^-|$. See Figure 5. In the later case, E intersects K in one point, and $V \cup E$ bounds a pair of a 3-ball and an unknotted subarc of K^- by the minimality of n . Then an isotopy of F along the pair can reduce $|F \cap B^-|$. See Figure 6. \square




FIGURE 5. Isotopy of F along the 3-ball




FIGURE 6. Isotopy of F along the pair

Lemma 3.4. *There is no face f of G in F such that ∂f is a loop of G .*

Proof. Suppose there exists a face f as Lemma 3.4. Then ∂f consists of an edge α of G and a subarc γ of the boundary of a vertex V of G . By Lemma 3.2, $p(\alpha)$ intersects $p(K^-)$. Moreover, since the loop $\gamma \cup p(\alpha)$ bounds a disk E in S , $|p(\alpha) \cap p(K^-)| = 1$ and γ meets exactly one of e_1 and e_2 , say e_1 . Thus a loop $l = \partial N(\partial E; E) - \partial E$ intersects \tilde{K} in two points. Since \tilde{K} is prime, $\text{int} E$ intersects $p(K)$ in an embedded arc. Then, there are two possibilities for e_1 , $e_1 \subset f$ or $e_1 \subset V$.

In the former case, $f \cup E$ bounds a pair of a 3-ball and an unknotted arc, and an isotopy of F along the pair eliminates α . In the latter case, $f \cup E$ bounds a 3-ball, and an isotopy of F along the 3-ball eliminates α . These contradict the minimality of $|(F \cap D^+) - (e_1 \cup e_2)|$. \square

Hence we have a condition that G has at least two vertices, every vertex has valency at least two, and all faces of G in F are disks. Next, we pay attention to a face of G in F .

Lemma 3.5. *For any face f , the cycle ∂F can not be oriented.*

Proof. If all corners of f do not meet $e_1 \cup e_2$, then this is same to Lemma 2.4.

If exactly one corner of f meets e_1 or e_2 at one point, then f and some K_i^+ have the intersection number ± 1 , or a vertex which meets f along the corner intersects some K_k^- in one point. Since $p(\partial f)$ and $p(K^-) \cap p(K_i^+)$ must have the intersection number zero, ∂f is bounded by a loop of G consisting of a vertex and an edge α , and $p(\alpha)$ intersects $p(K^-)$ in one point. Then Lemma 3.4 gives the conclusion.

If some corners of f meet both e_1 and e_2 , then the corners of f have the intersection number zero with $p(K)$ because F and K have the intersection number zero. In such a situation, we have a contradiction same as the proof of Lemma 2.4. \square

By Lemma 3.5, starting a face f of G in F whose closure is a disk, we can get a loop l in $F - K$ with $|lk(l, K)| \geq 2$. But this is impossible because any loop in $F - K$ is null-homotopic in $S^3 - K$ or has linking number ± 1 with K . This finishes the proof of Theorem 1.4.

REFERENCES

- [1] P. R. Cromwell, *Positive braids are visually prime*, Proc. London Math. Soc. (3) **67** (1993) 384-424.
- [2] M. Ozawa, *Synchronism of an incompressible non-free Seifert surface for a knot and an algebraically split closed surface in the knot complement*, Proc. Amer. Math. Soc. **128** (2000) 919-922.

DEPARTMENT OF MATHEMATICS, SCHOOL OF EDUCATION, WASEDA UNIVERSITY, NISHIWASEDA
1-6-1, SHINJUKU-KU, TOKYO 169-8050, JAPAN
E-mail address: ozawa@musubime.com